

A family of covering properties for forcing axioms and strongly compact cardinals

Matteo Viale
 KGRC, University of Vienna
 Equipe de Logique Mathématique, Université Paris 7
 matteo@logic.univie.ac.at

Abstract

This¹ paper presents the main results in my Ph.D. thesis and develops from [25] where it was shown that SCH follows from PFA. In what follows several other and simpler proofs of SCH are presented introducing a family of covering properties $CP(\kappa, \lambda)$ which implies both SCH and the failure of square. I will also apply these covering properties to investigate models of strongly compact cardinals or of strong forcing axioms like MM or PFA.

In this paper I introduce a family of covering properties $CP(\kappa, \lambda)$ indexed by pairs of regular cardinals $\lambda < \kappa$. In the first part I show that these covering properties capture the combinatorial content of many of the known proofs of the failure of square-like principles from forcing axioms or large cardinals. In the second part I show that a large class of these covering properties follows either from the existence of a strongly compact cardinal or from at least two combinatorial principles which hold under PFA and are mutually independent: the mapping reflection principle MRP introduced by Moore in [14] and the P -ideal dichotomy PID introduced in its full generality by Todorćević in [21] developing on preceding works by him and Abraham [1] on P -ideals of countable subsets of ω_1 . This allows for a unified and simple proof of the failure of square and of the singular cardinal hypothesis SCH assuming PFA. Finally in the last part of the paper I will investigate the "saturation properties"² of models of $CP(\kappa, \lambda)$ for various κ and λ .

The paper is organized as follows: in sections 1 and 2 we introduce the combinatorial principles PID and MRP. In sections 3 and 4 we introduce the covering properties $CP(\kappa, \lambda)$ and we outline some of their consequences among which SCH and the failure of square. In sections 5, 6 and 7 we prove various instances

¹This research has been partially supported by the Austrian Science Fund FWF project P19375-N18. I thank Boban Velickovic for many useful suggestions and comments on the redaction of this paper.

²In a sense that will be made explicit in the last section of the paper.

of $\text{CP}(\kappa, \lambda)$ assuming respectively the existence of a strongly compact cardinal, PID or MRP^3 . In the last section we study the rigidity of models of $\text{CP}(\kappa, \lambda)$ and we prove several results that strongly suggest that any two models $W \subseteq V$ of MM with the same cardinals have the same ω_1 -sequences of ordinals. Another result that we will present is that strongly compact cardinals λ are destroyed by any forcing which preserves cardinals and changes the cofinality of some regular $\kappa > \lambda$ to some $\theta < \lambda$.

The paper aims to be accessible and self-contained for any reader with a strong background in combinatorial set theory. Moreover while forcing axioms are a source of inspiration for the results that we present, all the technical arguments in this paper (except some of those in the last section) can be followed by a reader with no familiarity with the forcing techniques. When not otherwise explicitly stated [7] is the standard source for notation and definitions. For a regular cardinal θ , we use $H(\theta)$ to denote the structure $\langle H(\theta), \in, < \rangle$ whose domain is the collection of sets whose transitive closure is of size less than θ and where $<$ is a predicate for a fixed well ordering of $H(\theta)$. For cardinals $\kappa \geq \lambda$ we let $[\kappa]^\lambda$ be the family of subsets of κ of size λ . In a similar fashion we define $[\kappa]^{<\lambda}$, $[\kappa]^{\leq \lambda}$, $[X]^\lambda$, where X is an arbitrary set. If X is an uncountable set and θ a regular cardinal, $\mathcal{E} \subseteq [X]^\theta$ is unbounded if for every $Z \in [X]^\theta$, there is $Y \in \mathcal{E}$ containing Z . $\mathcal{E}$ is bounded otherwise. $\mathcal{E}$ is closed in $[X]^\omega$ if whenever $X = \bigcup_n X_n$ and $X_n \subseteq X_{n+1}$ are in $\mathcal{E}$ for all n , then also $X \in \mathcal{E}$. It is a well known fact that $\mathcal{C} \subseteq [X]^\omega$ is closed and unbounded (club) iff there is $f : [X]^{<\omega} \rightarrow X$ such that $\mathcal{C}$ contains the set of all $Y \in [X]^\omega$ such that $f[Y]^{<\omega} \subseteq Y$. $S \subseteq [X]^\omega$ is stationary if it intersects all club subsets of $[X]^\omega$. The f -closure of X is the smallest Y containing X such that $f[[Y]^{<\omega}] \subseteq Y$. Given f as above, $\mathcal{E}_f$ is the club of $Z \in [X]^\omega$ such that Z is f -closed. If X is a set of ordinals, then $\overline{X}$ denotes the topological closure of X in the order topology. For regular cardinals $\lambda < \kappa$, $S_\kappa^{\leq \lambda}$ denotes the subset of κ of points of cofinality $\leq \lambda$. In a similar fashion we define S_κ^λ and $S_\kappa^{<\lambda}$. We say that a family $\mathcal{D}$ is covered by a family $\mathcal{E}$ if for every $X \in \mathcal{D}$ there is a $Y \in \mathcal{E}$ such that $X \subseteq Y$. We also recall the following definitions central to the arguments that follows:

The singular cardinal hypothesis SCH asserts that $\kappa^{\text{cof}(\kappa)} = \kappa^+ + 2^{\text{cof}(\kappa)}$ for all infinite cardinals κ .

It is a celebrated result of Silver [19] that if SCH fails, then it first fails at a singular cardinal of countable cofinality. It is also known that the failure of SCH is a strong hypothesis as it entails the existence of models of ZFC with measurable cardinals [6].

Let κ be an infinite regular cardinal. The square principle $\square(\kappa)$ asserts the existence of a sequence $(C_\alpha : \alpha < \kappa)$ with the following properties:

- (i) for every limit α , C_α is a closed unbounded subset of α ,
- (ii) if α is a limit point of C_β , $C_\alpha = C_\beta \cap \alpha$,

³The shortest path to obtain a self-contained proof of the SCH starting from PID is to read sections 1, 3 and 6, from MRP is to read sections 2, 3 and 7.

- (iii) there is no club C in κ such that for all α there is $\beta \geq \alpha$ such that $C \cap \alpha = C_\beta$,
- (iv) $C_{\beta+1} = \{\beta\}$.

It is well known that the failure of $\square(\kappa)$ for all cardinals $\kappa > \aleph_1$ is a very strong large cardinal hypothesis. For example Schimmerling has shown that it entails the existence of models of set theory with Woodin cardinals [17].

Recall that λ is a strongly compact cardinal if for every $\kappa \geq \lambda$ there is a λ -complete ultrafilter on $[\kappa]^{<\lambda}$.

1 The P -ideal dichotomy

Let Z be an uncountable set. $\mathcal{I} \subseteq [Z]^{\leq \omega}$ is a P -ideal if it is an ideal and for every countable family $\{X_n\}_n \subseteq \mathcal{I}$ there is an $X \in \mathcal{I}$ such that for all n , $X_n \subseteq^* X$ (where $\subseteq^*$ is inclusion modulo finite).

Definition 1 (*Todorćević, [21]*)

The P -ideal dichotomy (PID) asserts that for every P -ideal $\mathcal{I}$ on $[Z]^{\leq \omega}$ for some fixed uncountable Z , one of the following holds:

- (i) *There is Y uncountable subset of Z such that $[Y]^{\leq \omega} \subseteq \mathcal{I}$.*
- (ii) *$Z = \bigcup_n A_n$ with the property that A_n is orthogonal to $\mathcal{I}$ (i.e. $X \cap Y$ is finite for all $X \in [A_n]^\omega$ and $Y \in \mathcal{I}$) for all n .*

PID is a principle which follows from PFA and which is strong enough to rule out many of the standard consequences of $V = L$. For example Abraham and Todorćević [1] have shown that under PID there are no Souslin trees while Todorćević has shown that PID implies the failure of $\square(\kappa)$ on all regular $\kappa > \aleph_1$ [21]. Due to this latter fact the consistency strength of this principle is considerable. Another interesting result by Todorćević is that PID implies that $\mathfrak{b} \leq \aleph_2^4$. Nonetheless in [1] and [21] it is shown that this principle is consistent with CH. Other interesting applications of PID can be found in [24], [2], [21] and [1].

2 The mapping reflection principle

Almost all known applications of MM which do not follow from PFA are a consequence of some form of reflection for stationary sets. These types of reflection principles are a fundamental source in order to obtain proofs of all cardinal arithmetic result that follows from MM. In particular SCH and the fact that

⁴In [21] it is shown that any gap in $P(\omega)/FIN$ is either an Hausdorff gap or a (κ, ω) gap with κ regular and uncountable. By another result of Todorćević (see [7] pp. 578 for a proof) if $\mathfrak{b} > \aleph_2$ there is an (ω_2, λ) gap in $P(\omega)/FIN$ for some regular uncountable λ . Thus PID is not compatible with $\mathfrak{b} > \aleph_2$.

$\mathfrak{c} \leq \omega_2$ are a consequence of many of the known reflection principles which hold under MM. However up to a very recent time there was no such kind of principle which could be derived from PFA alone. This has been the main difficulty in the search for a proof that PFA implies $\mathfrak{c} = \omega_2$, a result which has been obtained by Todorćević and Velićković appealing to combinatorial arguments which are not dissimilar from the P -ideal dichotomy [23]. Later on this has also been the crucial obstacle in the search for a proof of SCH from PFA.

In 2003 Moore [14] found an interesting form of reflection which can be derived from PFA, the mapping reflection principle MRP. He has then used this principle to show that BPFA implies that $\mathfrak{c} = \aleph_2$ and also that this principle is strong enough to entail the non-existence of square sequences. He has also shown in [16] that MRP could be a useful tool in the search of a proof of SCH from PFA. I first obtained my proof of this latter theorem elaborating from [16]. Many other interesting consequences of this reflection principle have been found by Moore and others. A complete presentation of this subject will be found in [4].

Definition 2 *Let θ be a regular cardinal, let X be uncountable, and let $M \prec H(\theta)$ be countable such that $[X]^\omega \in M$. A subset Σ of $[X]^\omega$ is M -stationary if for all $\mathcal{E} \in M$ such that $\mathcal{E} \subseteq [X]^\omega$ is club, $\Sigma \cap \mathcal{E} \cap M \neq \emptyset$.*

Recall that the Ellentuck topology on $[X]^\omega$ is obtained by declaring a set open if it is the union of sets of the form

$$[x, N] = \{Y \in [X]^\omega : x \subseteq Y \subseteq N\}$$

where $N \in [X]^\omega$ and $x \subseteq N$ is finite.

Definition 3 *Σ is an open stationary set mapping if there is an uncountable set X and a regular cardinal θ such that $[X]^\omega \in H(\theta)$, the domain of Σ is a club in $[H(\theta)]^\omega$ of countable elementary submodels M such that $X \in M$ and for all M , $\Sigma(M) \subseteq [X]^\omega$ is open in the Ellentuck topology on $[X]^\omega$ and M -stationary.*

The mapping reflection principle (MRP) asserts that:

If Σ is an open stationary set mapping, there is a continuous $\in$ -chain $\vec{N} = (N_\xi : \xi < \omega_1)$ of elements in the domain of Σ such that for all limit ordinals $0 < \xi < \omega_1$ there is $\nu < \xi$ such that $N_\eta \cap X \in \Sigma(N_\xi)$ for all η such that $\nu < \eta < \xi$.

If $(N_\xi : \xi < \omega_1)$ satisfies the conclusion of MRP for Σ then it is said to be a reflecting sequence for Σ .

We list below some of the interesting consequences of MRP.

- (Moore [14]) PFA implies MRP.
- (Moore [14]) MRP implies that $\mathfrak{c} = \aleph_2$. As a simple outcome of his proof of the above theorem Moore obtains also that BPFA implies $\mathfrak{c} = \aleph_2$.
- (Moore [14]) Assume MRP. Then $\square(\kappa)$ fails for all regular $\kappa \geq \aleph_2$.

A folklore problem in combinatorial set theory for the last twenty years has been the consistency of the existence of a five element basis for the uncountable linear orders, i.e. the statement that there are five uncountable linear orders such that at least one of them embeds in any other uncountable linear order.

- (Moore [15]) Assume BPFA and MRP. Then there is a five element basis for the uncountable linear orders.

A considerable reduction of the large cardinal hypothesis needed for the consistency of the above conjecture has been obtained in [10]. A byproduct of their results yields to the following:

- (König, Larson, Moore, Veličković [10]) MRP implies that there are no Kurepa trees.

Other interesting consequences of MRP can be found in [5].

We remark that MRP and PID are mutually independent principles since PID is compatible with CH while, by a result of Myiamoto [13], MRP is compatible with the existence of Souslin trees.

3 A family of covering properties

In this section we introduce the main original concept of this paper. A careful analysis of Todorčević's proof that PID implies that $\square(\kappa)$ fails for all regular $\kappa \geq \aleph_1$ leads to the isolation of a family of covering properties $\text{CP}(\kappa, \lambda)$ which on one side are strong enough to entail both SCH and the failure of square, and on the other side are weak enough to be a consequence of the existence of a strongly compact cardinal, of PID and of MRP.

Definition 4 For regular cardinals $\lambda < \kappa$, $\mathcal{D} = (K(\alpha, \beta) : \alpha < \lambda, \beta \in \kappa)$ is a λ -covering matrix for κ if:

- (i) $\beta \subseteq \bigcup_{\alpha < \lambda} K(\alpha, \beta)$ for all β ,
 - (ii) $|K(\alpha, \beta)| < \kappa$ for all β and α ,
 - (iii) $K(\alpha, \beta) \subseteq K(\eta, \beta)$ for all $\beta < \kappa$ and for all $\alpha < \eta < \lambda$,
 - (iv) for all $\gamma < \beta < \kappa$ and for all $\alpha < \lambda$, there is $\eta < \lambda$ such that $K(\alpha, \gamma) \subseteq K(\eta, \beta)$.
 - (v) for all $X \in [\kappa]^{\leq \lambda}$, there is $\gamma_X < \kappa$ such that for all $\beta < \kappa$ and $\eta < \lambda$, there is α such that $K(\eta, \beta) \cap X \subseteq K(\alpha, \gamma_X)$
- $\beta_{\mathcal{D}} \leq \kappa$ is the least β such that for all α and γ , $\text{otp}(K(\alpha, \gamma)) < \beta$

We will mainly be interested in ω -covering matrices, which we will just call covering matrices. As we will see below square like principles are useful to construct several kinds of covering matrices. One successful strategy to negate square principles from large cardinals and forcing axioms is to use appropriate ultrafilters or specific forcing arguments to "diagonalize" through the covering matrix defined appealing to these square-like principles, as for example in the proofs of the failure of square from a strongly compact by Solovay [20] or from PID by Todorćević [21]⁵. The covering matrices induced by square-like principles that we will consider satisfy a stronger coherence property than the "local" property (v). This condition is replaced by the "global" property⁶:

(v') For all $\gamma < \beta < \kappa$ and $\eta < \lambda$, there is α such that $K(\eta, \beta) \cap \gamma \subseteq K(\alpha, \gamma)$.

The key point to introduce condition (v) above in this weak form is that λ -covering matrices on κ can be defined in an elementary way in ZFC and the diagonalization argument which in the square-like cases leads to a contradiction, in the general case leads to a simple combinatorial argument to compute κ^λ . It is possible to prove the following:

Lemma 5 Assume that there is a stationary set of points of uncountable cofinality in the approachability ideal⁷ $\mathcal{I}[\kappa]$. Then there is an ω -covering matrix on κ . Moreover if λ is singular of countable cofinality, then there is $\mathcal{D}$ covering matrix on λ^+ with $\beta_{\mathcal{D}} = \lambda$.

Since the main original application of the existence of ω -covering matrices $\mathcal{D}$ that we have found is the proof of SCH from PFA, we will just prove a weaker form of the lemma:

Lemma 6 Assume that $\kappa > \mathfrak{c}$ is a singular cardinal of countable cofinality. Then there is an ω -covering matrix $\mathcal{C}$ for κ^+ with $\beta_{\mathcal{C}} = \kappa$.

Proof: The matrices we are going to define satisfy (i), ..., (iii) and a stronger coherence property than what is required by (iv) and (v) of the above definition. They will satisfy the following properties (iv*) and (v*) from which (iv) and (v) immediately follow:

(iv*) For all $\alpha < \beta$ there is n such that $K(m, \alpha) \subseteq K(m, \beta)$ for all $m \geq n$.

(v*) For all $X \in [\kappa^+]^\omega$ there is $\gamma_X < \kappa$ such that for all $\beta \geq \gamma_X$ there is n such that $K(m, \beta) \cap X = K(m, \gamma_X) \cap X$ for all $m \geq n$.

⁵See also the several arguments of this sort appearing in the sequel of this paper.

⁶For example the matrix produced by a square sequence using the ρ_2 -function (see sections 6 and 8 of [22] and fact 20 below) satisfies (i), ..., (iv) and (v'), and this matrix can be used to show that CP implies the failure of square. Another interesting example of a covering matrix which is not defined appealing to lemma 6 below and which satisfies (v) is the matrix used in the proof of theorem 27 in the last section.

⁷In [18] it is possible to find a definition of the approachability ideal $\mathcal{I}[\kappa]$. We avoid it in this paper since it is not relevant for the arguments we are presenting.

Let $\phi_\eta : \kappa \rightarrow \eta$ be a surjection for all $0 < \eta < \kappa^+$. Fix also $\{\kappa_n : n < \omega\}$ increasing sequence of regular cardinals cofinal in κ with $\kappa_0 \geq \aleph_1$. Set

$$K(n, \beta) = \bigcup \{K(n, \gamma) : \gamma \in \phi_\beta[\kappa_n]\}.$$

It is immediate to check that $\mathcal{D} = (K(n, \beta) : n \in \omega, \beta < \kappa^+)$ satisfies (i), (ii), (iii) of definition 4, property (iv*) and that $\beta_{\mathcal{D}} = \kappa$. To prove (v*), let $X \in [\kappa^+]^\omega$ be arbitrary. Now since $\mathfrak{c} < \kappa^+$ and there are at most $\mathfrak{c}$ many subsets of X , there is a stationary subset S of κ^+ and a fixed decomposition of X as the increasing union of sets X_n such that $X \cap K(n, \alpha) = X_n$ for all α in S and for all n . Now properties (i) $\dots$ (iv) of the matrix guarantees that this property of S is enough to get (v*) for X with $\gamma_X = \min(S)$. ■

A similar argument can be used to prove the following:

Lemma 7 *Assume that κ and λ are regular with $\kappa > 2^\lambda$. Then there is a λ -covering matrix $\mathcal{C}$ for κ .*

We say that $\mathcal{D}$ is covered by $\mathcal{E}$ iff every $X \in \mathcal{D}$ is contained in some $Y \in \mathcal{E}$.

Definition 8 $\text{CP}(\kappa, \lambda)$: κ has the λ -covering property⁸ if for every $\mathcal{D}$, λ -covering matrix for κ there is an unbounded subset A of κ such that $[A]^\lambda$ is covered by $\mathcal{D}$. $\text{CP}(\kappa)$ abbreviates $\text{CP}(\kappa, \omega)$ and CP is the statement that $\text{CP}(\kappa)$ holds for all regular $\kappa > \mathfrak{c}$.

Fact 9 *Assume $\text{CP}(\kappa^+)$ for all singular κ of countable cofinality. Then $\lambda^{\aleph_0} = \lambda$, for every $\lambda \geq 2^{\aleph_0}$ of uncountable cofinality.*

Proof: By induction. The base case is trivial. If $\lambda = \kappa^+$ with $\text{cof}(\kappa) > \omega$, then $\lambda^{\aleph_0} = \lambda \cdot \kappa^{\aleph_0} = \lambda \cdot \kappa = \lambda$, by the inductive hypothesis on κ . If λ is a limit cardinal and $\text{cof}(\lambda) > \omega$ then $\lambda^{\aleph_0} = \sup\{\mu^{\aleph_0} : \mu < \lambda\}$, so the result also follows by the inductive hypothesis. Thus, the only interesting case is when $\lambda = \kappa^+$, with κ singular of countable cofinality. In this case we will show, using CP , that $(\kappa^+)^{\aleph_0} = \kappa^+$. To this aim let $\mathcal{D}$ be a covering matrix for κ^+ with $\beta_{\mathcal{D}} = \kappa$. Notice that by our inductive assumptions, since every $K(n, \beta)$ has order type less than κ , $|[K(n, \beta)]^\omega|$ has size less than κ . So $\bigcup\{[K(n, \beta)]^\omega : n < \omega \text{ \& } \beta \in \kappa^+\}$ has size κ^+ . Use CP to find $A \subseteq \kappa^+$ unbounded in κ^+ , such that $[A]^\omega$ is covered by $\mathcal{D}$. Then $[A]^\omega \subseteq \bigcup\{[K(n, \beta)]^\omega : n < \omega \text{ \& } \beta \in \kappa^+\}$, from which the conclusion follows. ■

The following theorems motivate the introduction of these covering properties:

Theorem 10 *Assume λ is strongly compact. Then $\text{CP}(\kappa, \theta)$ holds for all regular $\theta < \lambda$ and all regular $\kappa \geq \lambda$.*

Theorem 11 *Assume PID. Then CP holds.*

⁸Moore has first noticed that a covering property similar to the ω -covering property for a regular $\kappa > \mathfrak{c}$ followed from MRP , reading a draft of [25].

On the other hand MRP allows us to infer a slightly weaker conclusion than the one of the previous theorem.

Theorem 12 *Assume MRP and let $\mathcal{D}$ be a covering matrix for κ such that $K(n, \beta)$ is a closed set of ordinals for all $K(n, \beta)$. Then there is A unbounded in κ such that $[A]^\omega$ is covered by $\mathcal{D}$.*

In particular we obtain:

Corollary 13 *PFA implies SCH.*

Proof: PFA implies PID and PID implies CP. In particular PFA implies that $\kappa^\omega = \kappa$ for all regular $\kappa \geq \mathfrak{c}$. By Silver's theorem [19] the least singular $\kappa > 2^{\text{cof} \kappa}$ such that $\kappa^{\text{cof} \kappa} > \kappa^+$ has countable cofinality. Now assume PFA and let κ have countable cofinality. By fact 9, $\kappa^{\text{cof}(\kappa)} \leq (\kappa^+)^{\aleph_0} = \kappa^+$. Thus assuming PFA there cannot be a singular cardinal of countable cofinality which violates SCH. Combining this fact with Silver's result we get that SCH holds under PFA. ■

Before proving all the above theorems we analyze in more details the effects of CP and we give other interesting examples of λ -covering matrices.

4 Some simple consequences of $\text{CP}(\kappa, \lambda)$

We remind the reader that if $\lambda^{\aleph_0} = \lambda$ for all regular $\lambda > 2^{\aleph_0}$, then by Silver's theorem and some elementary cardinal arithmetic we obtain that $\kappa^\lambda = \kappa + 2^\lambda$ for all regular cardinals $\lambda < \kappa$.

Fact 14 *Let $\theta < \kappa$ be regular cardinals such that $\lambda^\theta < \kappa$ for all regular $\lambda < \kappa$. $\mathcal{D} = \{K(\alpha, \beta) : \alpha \in \theta, \beta < \kappa\}$ be a θ -covering matrix on κ and A be an unbounded subset of κ . The following are equivalent:*

- (i) $[A]^\theta$ is covered by $\mathcal{D}$.
- (ii) $[A]^\lambda$ is covered by $\mathcal{D}$ for all $\lambda < \kappa$ such that $\text{cof}(\lambda) > \theta$.

Proof: (ii) implies (i) is evident. To prove the other direction, assume (i) and let $Z \subseteq A$ have size λ with $\text{cof}(\lambda) > \theta$. We need to find $\alpha < \theta$ and $\beta < \kappa$ such that $Z \subseteq K(\alpha, \beta)$. For $X \in [Z]^\theta \subseteq [A]^\theta$ let by (i) $\alpha_X < \theta$, $\beta_X < \kappa$ be such that $X \subseteq K(\alpha_X, \beta_X)$. By our assumptions, $\lambda^\theta < \kappa$. For this reason $\beta = \sup_{X \in [Z]^\theta} \beta_X < \kappa$. Now by property (iii) of $\mathcal{D}$, we have that for all $X \in [Z]^\theta$, $X \subseteq K(\alpha_X, \beta)$ for some α_X . Let $\mathcal{C}_\alpha$ be the set of X such that $\alpha_X = \alpha$. Now notice that for at least one α , $\mathcal{C}_\alpha$ must be unbounded in $[Z]^\theta$, otherwise $[Z]^\theta$ would be the union of θ bounded subsets which is not possible since $|Z|$ has cofinality larger than θ . Then $Z \subseteq K(\alpha, \beta)$, since every $\alpha \in Z$ is in some $X \in \mathcal{C}_\alpha$ because $\mathcal{C}_\alpha$ is unbounded. This completes the proof of the fact. □

A λ -covering matrix $\mathcal{D}$ for κ^+ with $\beta_{\mathcal{D}} = \kappa$ is an object simple to define when κ has cofinality λ . If $\kappa^\lambda = \kappa$, the existence of a covering matrix $\mathcal{D}$ for κ^+ with $\beta_{\mathcal{D}} < \kappa^+$ is not compatible with $\text{CP}(\kappa^+, \lambda)$. This is a simple consequence of the above facts:

Fact 15 *Assume $\kappa^\lambda = \kappa$ and $\text{CP}(\kappa^+, \lambda)$. Then there is no λ -covering matrix $\mathcal{D}$ on κ^+ with $\beta_{\mathcal{D}} < \kappa^+$.*

Proof: Assume not and let $\mathcal{D}$ be a λ -covering matrix for κ^+ with $\beta_{\mathcal{D}} < \kappa^+$. By $\text{CP}(\kappa^+, \lambda)$ there should be an A unbounded in κ^+ such that $[A]^\lambda$ is covered by $\mathcal{D}$. Appealing to fact 14, we can conclude in any case that $[A]^\kappa$ is covered by $\mathcal{D}$. Take β large enough in order that $\text{otp}(A \cap \beta) > \beta_{\mathcal{D}}$. Since $A \cap \beta$ has size at most κ there are n, γ such that $A \cap \beta \subseteq K(n, \gamma)$. Thus $\beta_{\mathcal{D}} < \text{otp}(A \cap \beta) \leq \text{otp}K(n, \beta) < \beta_{\mathcal{D}}$ a contradiction. ■

On the other hand assuming that κ is regular to what extent can we stretch λ in order that $\text{CP}(\kappa^+, \lambda)$ is consistent? The fact below shows that λ cannot be κ .

Lemma 16 *$\text{CP}(\kappa^+, \kappa)$ fails for all regular $\kappa \geq \omega_1$.*

Proof: Fix a sequence $\mathcal{C} = \{C_\xi : \xi < \kappa^+\}$ such that for all limit α , C_α is a club subset of α of order type $\text{cof}(\alpha)$. If $\xi = \alpha + 1$, $C_\xi = \{\alpha\}$. Define by induction on $\alpha \leq \beta < \kappa^+$,

$$\rho^*(\alpha, \beta) : [\kappa^+]^2 \rightarrow \kappa$$

as follows:

- $\rho^*(\alpha, \alpha) = 0$,
- $\rho^*(\alpha, \beta) = \max\{\text{otp}(C_\beta \cap \alpha), \rho^*(\alpha, \min(C_\beta \setminus \alpha)), \sup\{\rho^*(\xi, \alpha) : \xi \in C_\beta \cap \alpha\}\}$.

We will need the following properties of ρ^* which follows from the fact that κ is regular⁹.

Lemma 17 *For all $\alpha \leq \beta \leq \gamma$:*

- (a) $\rho^*(\alpha, \beta) \leq \max\{\rho^*(\alpha, \gamma), \rho^*(\beta, \gamma)\}$
- (b) $\rho^*(\alpha, \gamma) \leq \max\{\rho^*(\alpha, \beta), \rho^*(\beta, \gamma)\}$

□

Lemma 18 *For all $\alpha < \kappa^+$ and $\nu < \kappa$:*

$$|\{\xi < \alpha : \rho^*(\xi, \alpha) < \nu\}| \leq |\nu| + \aleph_0.$$

□

⁹For a proof see [22] lemmas 19.1 and 19.2

Set $D(\alpha, \beta) = \{\xi < \beta : \rho^*(\xi, \beta) \leq \alpha\}$ for all $\alpha < \kappa$ and $\beta < \kappa^+$.

Fact 19 *The following holds:*

- (i) $\text{otp}(D(\alpha, \beta)) < \kappa$ for all $\alpha < \kappa$ and $\beta < \kappa^+$,
- (ii) for all $\gamma < \beta < \kappa^+$, there is $\alpha_0 < \kappa$ such that $D(\alpha, \gamma) = D(\alpha, \beta) \cap \gamma$ for all $\alpha \geq \alpha_0$.

Proof: (i) follows from the second lemma on ρ^* . To prove (ii), let α_0 be such that $\rho^*(\gamma, \beta) \leq \alpha_0$, $\alpha \geq \alpha_0$ and $\xi < \beta$ such that $\rho^*(\xi, \beta) \leq \alpha$. By (a) of lemma 17:

$$\rho^*(\xi, \gamma) \leq \max\{\rho^*(\xi, \beta), \rho^*(\gamma, \beta)\} \leq \max\{\alpha, \alpha_0\} = \alpha.$$

Conversely assume that $\rho^*(\xi, \gamma) \leq \alpha$, by (b) of the same lemma:

$$\rho^*(\xi, \beta) \leq \max\{\rho^*(\xi, \gamma), \rho^*(\gamma, \beta)\} \leq \max\{\alpha, \alpha_0\} = \alpha.$$

Thus $D(\alpha, \gamma) = D(\alpha, \beta) \cap \gamma$ for all $\alpha \geq \alpha_0$. $\square$

This means that $\mathcal{D} = (D(\alpha, \beta) : \alpha < \kappa, \beta < \kappa^+)$ is a κ -covering matrix for κ^+ with $\beta_{\mathcal{D}} = \kappa$. Assuming $\text{CP}(\kappa^+, \kappa)$ there would be A unbounded in κ^+ such that $[A]^\kappa$ is covered by $\mathcal{D}$. However there cannot be an unbounded subset A of κ^+ such that $[A]^\kappa$ is covered by $\mathcal{D}$, since any element of the matrix has order type less than κ . $\square$

The function ρ^* defined above will be useful also for some other applications of $\text{CP}(\kappa, \lambda)$ in the final section (see theorem 27).

The following theorem follows closely Todorćević's proof that PID entails the failure of square and shows that CP is a very large cardinal property.

Theorem 20 *Assume $\kappa > \aleph_1$ is regular. Then $\text{CP}(\kappa)$ implies that $\square(\kappa)$ fails.*

Proof: Todorćević has shown that assuming $\square(\kappa)$ it is possible to define a step function (see sections 6 and 8 of [22]):

$$\rho_2 : [\kappa]^2 \rightarrow \omega$$

with the following properties:

- (i) For every A unbounded in κ , $\rho_2[[A]^2]$ is unbounded in ω ,
- (ii) for every $\alpha < \beta$ there is m such that $|\rho_2(\xi, \alpha) - \rho_2(\xi, \beta)| \leq m$ for all $\xi < \alpha$.

By (ii) it is immediate to check that $\mathcal{D} = \{K(n, \alpha) : n \in \omega \& \alpha < \kappa\}$ is a covering matrix for κ , where $K(n, \alpha) = \{\xi : \rho_2(\xi, \alpha) \leq n\}$. In fact it can be shown something stronger i.e. that for every $\alpha < \beta$ and n there is m such that $K(n, \alpha) \subseteq K(m, \beta)$ and $K(n, \beta) \cap \alpha \subseteq K(m, \alpha)$.

Using this coherence property of $\mathcal{D}$ one gets that whenever A is an unbounded subset of κ such that $[A]^\omega$ is covered by $\mathcal{D}$, then for all $\beta < \kappa$, $A \cap \beta \subseteq K(m_\beta, \beta)$ for some m_β . Thus one can refine any such A to an unbounded B such that for

a fixed m , $B \cap \beta \subseteq K(m, \beta)$ for all $\beta \in B$. This contradicts property (i) of ρ_2 . Assuming $\text{CP}(\kappa)$ we would get that an A unbounded in κ and such that $[A]^\omega$ is covered by $\mathcal{D}$ exists. However we just remarked that this is impossible. $\square$

The main difficulty towards a proof that PFA implies SCH has been the fact that all standard principles of reflection for stationary sets do not hold for PFA. In particular Baudoin [3] and Magidor (unpublished) have shown that PFA is compatible with the existence on any regular $\kappa \geq \aleph_2$ of a never reflecting stationary subset of S_κ^ω . However the following form of reflection holds under $\text{CP}(\kappa)$:

Fact 21 *Assume $\text{CP}(\kappa)$ for a regular $\kappa > \mathfrak{c}$ and let $\mathcal{D}$ be a covering matrix for κ with all $K(n, \beta)$ closed. Let $\lambda < \kappa$ be a regular cardinal and let $(S_\eta : \eta < \lambda)$ be an arbitrary family of stationary subsets of $S_\kappa^{\leq \lambda}$. Then there exist n and β such that $S_\eta \cap K(n, \beta)$ is non-empty for all $\eta < \lambda$.*

Proof: By $\text{CP}(\kappa)$ and facts 14, there is X unbounded in κ such that $[X]^\lambda$ is covered by $\mathcal{D}$. Since $K(n, \beta)$ is closed for all n and β , we have that $[\overline{X} \cap S_\kappa^{\leq \lambda}]^\lambda$ is covered by $\mathcal{D}$. To see this, let Z be in this latter set and find $Y \subseteq X$ of size λ such that $Z \subseteq \overline{Y}$. Now find n and β such that $Y \subseteq K(n, \beta)$. Since $K(n, \beta)$ is closed, $Z \subseteq \overline{Y} \subseteq K(n, \beta)$.

Now pick $M \prec H(\Theta)$ with Θ large enough such that $|M| = \lambda \subseteq M$ and $\lambda, X, (S_\eta : \eta < \lambda) \in M$. Then $S_\eta \cap \overline{X} \cap S_\kappa^{\leq \lambda}$ is non-empty for all η . By elementarity, M sees this and so $M \cap S_\eta \cap \overline{X} \cap S_\kappa^{\leq \lambda}$ is non-empty for all η . However $M \cap \overline{X} \cap S_\kappa^{\leq \lambda}$ has size λ so there are n and β such that $M \cap \overline{X} \cap S_\kappa^{\leq \lambda} \subseteq K(n, \beta)$. So $S_\eta \cap K(n, \beta)$ is non-empty for all η . $\square$

5 Strongly compact cardinals and $\text{CP}(\kappa, \theta)$

We turn to the proof of theorem 10. We will need the following trivial consequence of the existence of a strongly compact cardinal:

Lemma 22 *Assume λ is strongly compact. Then for every regular $\kappa \geq \lambda$, there is $\mathcal{U}$, λ -complete uniform ultrafilter on κ which concentrates on $S_\kappa^{< \lambda}$.*

Proof: Assume λ is strongly compact and $\kappa \geq \lambda$ is regular. By definition of λ there is a λ -complete ultrafilter $\mathcal{W}$ on $[\kappa]^{< \lambda}$ such that for all $X \in [\kappa]^{< \lambda}$, $\{Y \in [\kappa]^{< \lambda} : X \subseteq Y\} \in \mathcal{W}$. Set $\mathcal{U}$ to be the family of $A \subseteq \kappa$ such that $\{X \in [\kappa]^{< \lambda} : \sup(X \cap \alpha) = \alpha\} \in \mathcal{W}$ for all $\alpha \in A$. It is immediate to check that $\mathcal{U}$ is a λ -complete ultrafilter which concentrates on $S_\kappa^{< \lambda}$. $\square$

Now let $\theta < \lambda$ and $\kappa \geq \lambda$ be regular cardinals and fix a θ -covering matrix $\mathcal{D} = (K(\alpha, \beta) : \alpha \in \theta, \beta \in \kappa)$ for κ . Let $A_\alpha^\gamma = \{\beta > \gamma : \gamma \in K(\alpha, \beta)\}$ and $A_\alpha = \{\gamma \in S_\kappa^{< \lambda} : A_\alpha^\gamma \in \mathcal{U}\}$. Since $\theta < \lambda$, by the λ -completeness of $\mathcal{U}$, for every $\gamma \in S_\kappa^{< \lambda}$, there is a least α such that $A_\alpha^\gamma \in \mathcal{U}$. Thus $\bigcup_{\alpha < \theta} A_\alpha = S_\kappa^{< \lambda}$. So there is $\alpha < \theta$ such that $A_\alpha \in \mathcal{U}$. In particular A_α is unbounded. Now let X be a subset of A_α of size θ . Then $A_\alpha^\gamma \in \mathcal{U}$ for all $\gamma \in X$. Since $|X| = \theta < \lambda$, $\bigcap_{\gamma \in X} A_\alpha^\gamma \in \mathcal{U}$

and thus is non-empty. Pick β in this latter set. Then $X \subseteq K(\alpha, \beta)$. Since X is an arbitrary subset of A_α of size θ , we conclude that $[A_\alpha]^\theta$ is covered by $\mathcal{D}$. This concludes the proof of theorem¹⁰ 10. ■

6 PID implies CP

We turn to the proof of theorem 11. As we will see below a model of PID retains enough properties of the supercompact cardinals from which it is obtained in order that a variation of the above argument can be run also in the context of ω -covering matrices. We break the proof of theorem 11 in two parts. Assume κ is regular and let $\mathcal{D} = (K(n, \alpha) : n \in \omega, \alpha < \kappa)$ be a covering matrix on κ . Let $\mathcal{I}$ be the family of $X \in [\kappa]^\omega$ such that $X \cap K(n, \alpha)$ is finite for all $\alpha < \kappa$ and for all $n < \omega$.

Claim 23 $\mathcal{I}$ is a P -ideal.

Proof: Let $\{X_n : n \in \omega\} \subseteq \mathcal{I}$. Let $Y = \bigcup_n X_n$. Let γ_Y witness (v) for $\mathcal{D}$ relative to Y . Now since for every n, m , $X_n \cap K(m, \gamma_Y)$ is finite, let $X(n, m)$ be the finite set

$$X_n \cap K(m, \gamma_Y) \setminus K(m-1, \gamma_Y)$$

and let:

$$X = \bigcup_n \bigcup_{j \geq n} X(n, j).$$

Notice that $X_n = \bigcup_j X(n, j)$ and $\bigcup_{j \geq n} X(n, j) \subseteq X$, so we have that $X_n \subseteq^* X$. Moreover $X \cap K(n, \gamma_Y) = \bigcup_{j \leq i \leq n} X(j, i)$, so it is finite. We claim that $X \in \mathcal{I}$. If not there would be some β and some l such that $X \cap K(l, \beta)$ is infinite. Now $X \cap K(l, \beta) \subseteq Y \cap K(l, \beta) \subseteq K(m, \gamma_Y)$ for some m . Thus we would get that $X \cap K(m, \gamma_Y)$ is infinite for some m contradicting the very definition of X . □

Now notice that if $Z \subseteq \kappa$ is any set of ordinals of size $\aleph_1$ and $\alpha = \sup(Z)$, there must be an n such that $Z \cap K(n, \alpha)$ is uncountable. This means that $\mathcal{I} \not\subseteq [Z]^\omega$, since any countable subset of $Z \cap K(n, \alpha)$ is not in $\mathcal{I}$. This forbids $\mathcal{I}$ to satisfy the first alternative of the P -ideal dichotomy. So the second possibility must be the case, i.e. we can split κ in countably many sets A_n such that $\kappa = \bigcup_n A_n$ and for each n , $[A_n]^\omega \cap \mathcal{I} = \emptyset$.

Claim 24 For every n , $[A_n]^\omega$ is covered by $\mathcal{D}$.

Proof: Assume that this is not the case and let $X \in [A_n]^\omega$ be such that $X \setminus X(l, \beta)$ is non-empty for all l, β . Now let X_0 be a subset of X such that $X_0 \cap K(l, \gamma_X)$ is finite for all l . Then exactly as in the proof of claim 23 we can see that $X_0 \in [A_n]^\omega \cap \mathcal{I}$. This contradicts the definition of A_n . □

This concludes the proof of theorem 11. ■

¹⁰Notice that for the proof of this theorem we just needed property (i) of a covering matrix. The proof of $\mathbf{CP}(\kappa, \omega)$ assuming either $\mathbf{PID}$ or $\mathbf{MRP}$ will need properties (i) , (iii) , (iv) , (v) .

7 MRP implies SCH

We prove theorem 12. Thus assume MRP and let $\mathcal{D}$ be a covering matrix on a regular $\kappa > \mathfrak{c}$ such that $K(n, \beta)$ is a closed set of ordinals for all n and β . Assume that for all A unbounded in κ , $[A]^\omega$ is not covered by $\mathcal{D}$. We will reach a contradiction. For each $\delta < \kappa$ of countable cofinality, fix C_δ cofinal in δ of order type ω . Let M be a countable elementary submodel of $H(\Theta)$ for some large enough regular Θ . Let $\delta_M = \sup(M \cap \kappa)$ and β_M be the ordinal $\gamma_{M \cap \kappa}$ provided by property (v) of $\mathcal{D}$ applied to $M \cap \kappa$. Set $\Sigma(M)$ to be the set of all countable $X \subseteq M \cap \kappa$ bounded in δ_M such that

$$\sup(X) \notin K(|C_{\delta_M} \cap \sup(X)|, \beta_M).$$

We will show that $\Sigma(M)$ is open and M -stationary. Assume this is the case and let $\{M_\eta : \eta < \omega_1\}$ be a reflecting sequence for Σ . Let $\delta_{M_\xi} = \delta_\xi$ and $\delta = \sup_{\omega_1} \delta_\xi$. Find $C \subseteq \omega_1$ club such that $\{\delta_\xi : \xi \in C\} \subseteq K(n, \delta)$ for some n (which is possible since the $K(n, \delta)$ are closed subsets of κ). Let α be a limit point of C . Let $M = M_\alpha$ and notice that by our choice of β_M for all m , there is l such that $K(m, \delta) \cap M \subseteq K(l, \beta_M)$. This means that for all $\eta \in C \cap \alpha$, $\delta_\eta \in K(n, \delta) \cap M \subseteq K(l, \beta_M)$ for some fixed l . Since α is a limit point of C there is $\eta \in \alpha \cap C$ such that $|C_{\delta_M} \cap \delta_\eta| > l$ and $M_\eta \cap \kappa \in \Sigma(M)$. But this is impossible, since $M_\eta \cap \kappa \in \Sigma(M)$ means that $\delta_\eta \notin K(|C_{\delta_M} \cap \delta_\eta|, \beta_M)$, i.e. $\delta_\eta \notin K(l, \beta_M)$.

We now show that Σ_M is open and M -stationary:

Claim 25 $\Sigma(M)$ is open.

Proof: Assume $X \in \Sigma(M)$, we will find $\gamma \in X$ such that $[\{\gamma\}, X] \subseteq \Sigma(M)$. To this aim notice that $C_{\delta_M} \cap \sup(X)$ is a finite subset of X . Let $n_0 = |C_{\delta_M} \cap \sup(X)|$ and $\gamma_0 = \max(C_{\delta_M} \cap \sup(X)) + 1$. Since $X \in \Sigma(M)$, $\sup(X) \notin K(n_0, \beta_M)$ and so, since $K(n_0, \beta_M)$ is closed, $\gamma_1 = \max(K(n_0, \beta_M) \cap \sup(X)) < \sup(X)$. Thus, let $\gamma \in X$ be greater or equal than $\max\{\gamma_1 + 1, \gamma_0\}$. If $Y \in [\{\gamma\}, X]$, then $\gamma_0 \leq \sup(Y) \leq \sup(X)$, so $|C_{\delta_M} \cap \sup(Y)| = |C_{\delta_M} \cap \sup(X)| = n_0$ and

$$\gamma_1 = \max(K(n_0, \beta_M) \cap \sup(X)) < \sup(Y) \leq \sup(X) < \min(K(n_0, \beta_M) \setminus \sup(X)).$$

Thus $Y \notin K(|C_{\delta_M} \cap \sup(Y)|, \beta_M)$, i.e. $Y \in \Sigma(M)$. $\square$

Claim 26 $\Sigma(M)$ is M -stationary.

Proof: Let $f : [\kappa]^{<\omega} \rightarrow \kappa$ in M . We need to find $X \in \Sigma(M)$ such that $f[[X]^{<\omega}] = X$. Let $N \prec H(\kappa^+)$ be a countable submodel in M such that $f \in N$ and let $C = \{\delta < \kappa : f[[\delta]^{<\omega}] = \delta\}$. Let also $n_0 = |C_{\delta_M} \cap \sup(N \cap \kappa)|$ and $\gamma_0 \in N$ be larger than $\max(C_{\delta_M} \cap \sup(N \cap \kappa))$. Then $(C \setminus \gamma_0) \in N$. We assumed that no A unbounded in κ is such that $[A]^\omega$ is covered by $\mathcal{D}$. So in particular by elementarity of N :

$$N \models [(C \setminus \gamma_0) \cap S_\kappa^\omega]^\omega \text{ is not covered by } \mathcal{D}$$

Thus there exists $X \in N$ countable subset of $(C \setminus \gamma_0) \cap S_\kappa^\omega$ such that for all n and β , $X \setminus K(n, \beta)$ is non-empty. Let $\gamma \in X \setminus K(n_0, \beta_M)$. Now find $Z \in N$ countable and cofinal in γ and let Y be the f -closure of Z . Then $Y \in N \subseteq M$. Now $\gamma \in C$ so $\sup(Y) = \sup(Z) = \gamma \notin K(n_0, \beta_M)$. Moreover $\gamma = \sup(Y) \in (C \setminus \gamma_0) \cap N$, so $\gamma_0 < \sup(Y) < \sup(N \cap \kappa)$, i.e. $|C_{\delta_M} \cap \sup(Y)| = |C_{\delta_M} \cap \sup(N \cap \kappa)| = n_0$. Thus:

$$\sup(Y) \notin K(|C_{\delta_M} \cap \sup(Y)|, \beta_M).$$

I.e. $Y \in \Sigma(M)$. □

This concludes the proof of theorem 12. ■

8 "Saturation" properties of models of strong forcing axioms.

Since forcing axioms have been able to settle many of the classical problems of set theory, we can expect that the models of a forcing axiom are in some sense categorical. There are many ways in which one can give a precise formulation to this concept. For example, one can study what kind of forcing notions can preserve PFA or MM, or else if a model V of a forcing axiom can have an interesting inner model M of the same forcing axiom. There are many results in this area, some of them very recent. First of all there are results that shows that one has to demand a certain degree of resemblance between V and M . For example assuming large cardinals it is possible to use the stationary tower forcing introduced by Woodin¹¹ to produce two transitive models $M \subseteq V$ of PFA (or MM or whatever is not conflicting with large cardinal hypothesis) with different ω -sequences of ordinals and an elementary embedding between them. However M and V do not compute the same way neither the ordinals of countable cofinality nor the cardinals. On the other hand, König and Yoshinobu [11, Theorem 6.1] showed that PFA is preserved by ω_2 -closed forcing while it is a folklore result that MM is preserved by ω_2 -directed closed forcing. Notice however that all these forcing notion do not introduce new sets of size at most $\aleph_1$. In the other direction, in [23] Veličković used a result of Gitik to show that if MM holds and M is an inner model such that $\omega_2^M = \omega_2$, then $\mathcal{P}(\omega) \subseteq M$ and Caicedo and Veličković [5] showed, using the mapping reflection principle MRP introduced by Moore in [16], that if $M \subseteq V$ are models of BPFA and $\omega_2^M = \omega_2$ then $\mathcal{P}(\omega_1) \subseteq M$. In any case all the results so far produced show that any two models $V \subseteq W$ of some strong forcing axiom and with the same cardinals have the same ω_1 -sequences of ordinals. Thus it is tempting to conjecture that forcing axioms produce models of set theory which are "saturated" with respect to sets of size $\aleph_1$. One possible way to give a precise formulation to this idea may be the following:

Conjecture 1 (*Veličković*) *Assume $W \subseteq V$ are models of MM with the same cardinals. Then $[Ord]^{\leq \omega_1} \subseteq W$.*

¹¹[12] gives a complete presentation of this subject.

or in a weaker form:

Conjecture 2 *Assume $W \subseteq V$ are models of MM with the same cardinals. Then $[Ord]^\omega \subseteq W$.*

CP can be used to show that one shouldn't expect that the negation of conjecture 2 can be obtained by means of forcing. Since no inner model theory is known for models of MM, PFA or of a supercompact cardinal, the results I will present brings to the conclusion that currently there are no suitable means to try to prove the negation of the second conjecture. Moreover I can show that if this conjecture fails first at κ , then κ is close to be a Jónsson cardinal in the smaller model and more.

Theorem 27 *Assume $CP(\kappa^+, \theta)$. Let W be an inner model such that κ is a regular cardinal of W and such that $(\kappa^+)^W = \kappa^+$. Then $\text{cof}(\kappa) \neq \theta$.*

This shows that if λ is strongly compact, than one cannot change the cofinality of some regular $\kappa \geq \lambda$ to some $\theta < \lambda$ and preserve at the same time κ^+ and the strong-compactness of λ .

Corollary 28 *Assume PFA and let W be an inner model with the same cardinals. Then W computes correctly all the points of countable cofinality.*

Proof: PFA implies CP. Now apply the above theorem. ■

In particular this gives another proof that Prikry forcing destroys PFA, since if g is a Prikry generic sequence on a measurable κ , $V[g]$ cannot model CP.

The next propositions show that neither κ^+ -CC forcing notions nor homogeneous forcing notions can be used to obtain a generic extension which models PFA and which first adds a new ω -sequence to κ . First of all we can see that the failure of conjecture 2 entails that W and V have a diverging notion of stationarity:

Proposition 29 *Let W be an inner model of V and assume that V models CP. Assume that for the least W -cardinal κ such that $\kappa^\omega \setminus W$ is nonempty, $\kappa^+ = (\kappa^+)^W$ and $\kappa > \mathfrak{c}$. Moreover assume that $[\lambda]^\omega \subseteq W$ for all $\lambda < \kappa$. Then for every $\lambda < \kappa$ regular cardinal of V , there is S , W -stationary subset of $(S_{\kappa^+}^\lambda)^W$, which is not anymore stationary in V .*

In particular any forcing P satisfying the κ^+ -chain condition and such that κ is the least ordinal to which P adds a new ω -sequence destroys PFA, since these forcings preserves stationary subsets of κ^+ . Thus also diagonal Prikry forcing kills PFA.

Recall that $(P, <_P)$ is an homogeneous notion of forcing if for every $p, q \in P$, there are $p_0 \leq_P p$ and $q_0 \leq_P q$ such that $(\{r : r \leq p_0\}, <_P)$ is isomorphic to $(\{r : r \leq q_0\}, <_P)$. If P is an homogeneous notion of forcing it is possible to prove by a standard induction argument that if $\phi(a_0, \dots, a_n)$ is a formula in

the language of forcing whose parameters are canonical names for sets in the ground model, then either $\Vdash_P \phi(a_0, \dots, a_n)$ or $\Vdash_P \neg \phi(a_0, \dots, a_n)$. Notice that the standard cardinal preserving forcing to add an ω -sequence to an uncountable cardinal like Prikry forcing or diagonal Prikry forcing are homogeneous¹².

Proposition 30 *Let P be an homogeneous notion of forcing, let G be a P -generic filter. Assume that $\Vdash_P [\lambda]^\omega \subseteq V$ for all $\lambda < \kappa$ and that $\kappa^+ > \mathfrak{c}$ is preserved. Assume moreover that $V[G]$ models $\text{CP}(\kappa^+)$. Then $\Vdash_P [\kappa]^\omega \subseteq V$*

$\text{CP}(\kappa^+)$ follows both from PFA and from the existence of a supercompact $\lambda \leq \kappa$. In particular the above proposition shows that PFA, MM and supercompact cardinals are not preserved by all known forcing notions which add ω -sequences and preserve cardinals since all such forcing notions are homogeneous.

We can conclude that there are not much hopes to find a forcing which first falsify conjecture 2 at some κ . By theorem 27, κ must be of countable cofinality. Moreover we have to seek for a P which on one hand preserves cardinals and is stationary set preserving, on the other hand is not homogeneous and does not have the κ^+ -chain condition¹³.

The following propositions give some ideas on how to measure the consistency strength of the negation of the above conjectures.

Say that κ is $\mathcal{E}$ -Jónsson for a family $\mathcal{E}$ of functions $\psi : [\kappa]^{<\omega} \rightarrow \kappa$ if for every $\psi \in \mathcal{E}$ there is Y proper subset of κ which is ψ -closed. κ is Jónsson if $\mathcal{E}$ contains all the relevant functions.

Any measurable cardinal is Jónsson. Silver has shown¹⁴ that if $\aleph_\omega > \mathfrak{c}$ is Jónsson, then it is measurable in an inner model. It is a major open problem whether $\aleph_\omega$ can be a Jónsson cardinal in some model of ZFC and recently König [9] has shown that MM is consistent with $\aleph_\omega$ not being Jónsson.

Proposition 31 *Let $W \subseteq V$ be models of ZFC and $\kappa > \mathfrak{c}$ be the least such that $[\kappa]^\omega \setminus W$ is nonempty. Assume that $V \models \text{CP}$, $2^\omega < \kappa$ and that $(\kappa^+)^W = \kappa^+$. Then κ^+ is W -Jónsson.*

Proposition 32 *Let $W \subseteq V$ be models of ZFC with the same bounded subsets of κ and computing κ^+ the same way. Assume that $V \models \text{CP}$, $2^\omega < \kappa$, and that κ is the least such that $[\kappa]^\omega \not\subseteq W$. Then κ is Jónsson in W .*

We now turn to the proofs.

Proof of theorem 27: Assume the theorem is false. Then $\text{cof}(\kappa) = \theta$. We need some preparation. Work in W . Fix a sequence $\mathcal{C} = \{C_\xi : \xi < \kappa^+\} \in W$

¹²It should be expected that the stationary set preserving forcing used by Magidor to show that MM implies there is a stationary set of bad points in $\aleph_{\omega+1}$ is not homogeneous. In any case this forcing collapses $\aleph_{\omega+1}$ to an ordinal of size $\aleph_1$ and moreover CH holds in the extension.

¹³In fact more than that, since the proof of proposition 29 shows that this P should have the property of forcing the non-stationarity of at least one set in every disjoint family in the ground model of κ -many stationary (in the ground model) subsets of $S_{\kappa^+}^\lambda$.

¹⁴See [8] for a proof.

such that for all limit α , C_α is a club subset of α of order type $\text{cof}(\alpha)$. If $\xi = \alpha + 1$, $C_\xi = \{\alpha\}$. Consider the function $\rho^* : [\kappa^+]^2 \rightarrow \kappa$ defined from such a sequence which was introduced in the proof of lemma 16.

Now work in V . Let $g : \theta \rightarrow \kappa$ be a cofinal sequence. Set $D(\alpha, \beta) = \{\xi < \alpha : \rho^*(\xi, \beta) \leq g(\alpha)\}$. The two lemmas 17 and 18 allows to prove the analogue of fact 19:

Fact 33 *The following holds:*

- (i) $\text{otp}(D(\alpha, \beta)) < \kappa$ for all $\alpha < \theta$ and $\beta < \kappa^+$,
- (ii) for all $\gamma < \beta < \kappa^+$, there is $\alpha_0 < \theta$ such that $D(\alpha, \gamma) = D(\alpha, \beta) \cap \gamma$ for all $\alpha \geq \alpha_0$.

□

This means that $\mathcal{D} = (D(\alpha, \beta) : \alpha < \theta, \beta < \kappa^+)$ is a θ -covering matrix for κ^+ with $\beta_{\mathcal{D}} = \kappa$. By $\text{CP}(\kappa^+, \theta)$ there is A unbounded in κ^+ such that $[A]^\theta$ is covered by $\mathcal{D}$. We claim that $[A]^\kappa$ is covered by $\mathcal{D}$. Suppose not, then there would be some $\gamma \leq \beta$ such that $\text{otp}(A \cap \gamma) > \kappa$ and $A \cap \gamma \setminus D(\alpha, \beta)$ is non-empty for all $\alpha < \theta$. Thus we could find an $X \in [A \cap \gamma]^\theta$ such that $X \setminus D(\alpha, \beta)$ is non-empty for all $\alpha < \theta$. But then, by (ii) of the above fact, $X \setminus D(\alpha, \beta)$ would be non-empty for all $\alpha < \theta$ and $\beta < \kappa^+$. This would contradict the fact that $[A]^\theta$ is covered by $\mathcal{D}$. Now let γ be such that $\kappa < \text{otp}(A \cap \gamma)$. Then for some α and β we would have $A \cap \gamma \subseteq D(\alpha, \beta)$ and so:

$$\kappa < \text{otp}(A \cap \gamma) \leq \text{otp}(D(\alpha, \beta)) < \kappa.$$

This is the contradiction which proves the theorem. ■

Proof of proposition 29: By the previous theorem κ cannot be regular in W , otherwise V cannot model CP. Since κ is the least such that $\kappa^\omega \setminus W$ is non empty and is not regular in W , we can conclude that κ is in W a cardinal of countable cofinality. So there is a covering matrix $\mathcal{D} \in W$ for $(\kappa^+)^W$ with $\beta_{\mathcal{D}} = \kappa$ and such that $K(n, \beta)$ is closed for all n and β . Since $\kappa^+ = (\kappa^+)^W$, $\mathcal{D}$ is still a covering matrix in V for κ^+ with $\beta_{\mathcal{D}} = \kappa$. Assume towards a contradiction that there is some $\lambda < \kappa$ regular cardinal of V such that every W -stationary subset of $S_{\kappa^+}^\lambda$ is still stationary in V . So fix in W some $\{A_\alpha : \alpha < \kappa\} \in W$ partition of $(S_{\kappa^+}^\lambda)^W$ in κ -many W -stationary sets. Now by our assumption, this is still a family of disjoint stationary subsets of V . Let for every n, β , $D(n, \beta)$ be the set of $\alpha < \kappa$ such that $A_\alpha \cap K(n, \beta)$ is nonempty. Now $D(n, \beta) \in W$ and $|D(n, \beta)| \leq |K(n, \beta)|$ has size less than κ . So, by minimality of κ , we have that $D(n, \beta)^\omega \subseteq W$, else there would be a new ω -sequence in $|D(n, \beta)|$. Apply CP in V and find X unbounded in κ^+ such that $[X]^\omega$ is covered by $\mathcal{D}$. Exactly as in the proof of fact 21, we can see that

$$[\overline{X} \cap S_{\kappa^+}^{\leq \lambda}]^\lambda$$

is covered by $\mathcal{D}$. Now pick M countable elementary submodel containing all relevant information. Then $A_{g(n)} \cap \overline{X} \cap M \cap S_{\kappa^+}^\lambda$ is nonempty for all n , by elementarity of M . Now $M \cap \overline{X} \cap S_{\kappa^+}^\lambda \subseteq K(n, \beta)$ for some n, β . This means that $g \in D(n, \beta)^\omega \subseteq W$ and we are done. ■

Proof of proposition 30: As before we can remark that κ has countable cofinality in V , otherwise the hypothesis of the proposition are not compatible with the conclusion of theorem 27. For this reason $S_{\kappa^+}^\omega$ has the same interpretation in V and in $V[G]$. Now we will use homogeneity to obtain a family $T \in V$ of κ -many disjoint subsets of $S_{\kappa^+}^\omega$ which are stationary in $V[G]$, we will then proceed exactly as in the previous proposition to obtain the desired conclusion. To this aim let:

$$\{A_{\alpha\beta} : \alpha < \kappa, \beta \in \kappa^+\} \in V$$

be an Ulam matrix, i.e for all $\alpha < \kappa$ and $\beta < \kappa^+$,

$$A_{\alpha\beta} = \{\xi \in S_{\kappa^+}^\omega : \phi_\xi(\beta) = \alpha\}$$

where $\phi_\xi : \xi \rightarrow \kappa \in V$ is a bijection for all $\xi \in [\kappa, \kappa^+)$.

A standard ZFC argument shows that there is a row α of the matrix such that the set of β such that $A_{\alpha\beta}$ is stationary has size κ^+ , moreover $A_{\alpha\beta} \cap A_{\alpha\gamma} = \emptyset$ whenever $\beta \neq \gamma$.

Fix an α with this property in $V[G]$ and let S be the set of β such that $V[G]$ models $A_{\alpha\beta}$ is stationary. Now by the homogeneity of P and the fact that $A_{\alpha\beta} \in V$ we can conclude that:

$$\Vdash_P A_{\alpha\beta} \text{ is a stationary subset of } S_{\kappa^+}^\omega$$

for all $\beta \in S$. We can conclude that $S \in V$. Let $T = \{S_\alpha : \alpha < \kappa\} \in V$ be a subset of S of size κ . Now proceed exactly as in the previous proposition. ■

Proof of propositions 31 and 32: Remark that by theorem 27 any κ as in the hypothesis of theorems 31 and 32 has already countable cofinality in W . Fix $A \in V$ unbounded set witnessing CP relative to a $\mathcal{D} \in W$ covering matrix for κ^+ with $\beta_{\mathcal{D}} = \kappa$. Notice that this entails that $[A]^\omega \subseteq W$. Now let $\psi : [\kappa^+]^{<\omega} \rightarrow \kappa^+$ be any function in W . Let B be the ψ -closure of A . We claim that B is a proper subset of κ^+ , otherwise if $X \in [\kappa^+]^\omega \setminus W$ is contained in B there would be a countable subset Y of A such that X is contained in the ψ -closure Z of Y . However $[Z]^\omega \subseteq W$, since $Z \in W$ is countable, so $X \in W$ and this contradicts our choice of X . This shows that κ^+ is W -Jónsson. To show proposition 32, set for every $\psi : [\kappa]^{<\omega} \rightarrow \kappa$, $(T_\psi, <_{T_\psi})$ to be the well-founded partial order of bounded subsets X of κ such that X is strictly contained in $\sup(X)$ and such that for all $\alpha_1, \dots, \alpha_n \in X$, if $\psi(\alpha_1, \dots, \alpha_n) < \sup(X)$, then $\psi(\alpha_1, \dots, \alpha_n) \in X$. If $X, Y \in T_\psi$, $X <_{T_\psi} Y$ if Y end extends X and $|X| < |Y|$. Now notice that T_ψ gets the same interpretation in W and in V . If $\bar{\psi} : [\kappa^+]^{<\omega} \rightarrow \kappa^+$ is in W an arbitrary extension of ψ and A witness in V the covering property, $\bar{\psi}[A] \cap \kappa$ is a branch through T_ψ of type κ . Thus V models that $>_{T_\psi}$ is ill-founded. By absoluteness W models that $>_{T_\psi}$ is ill-founded.

Notice that our requirement on the order $<_{T_\psi}$ entails that a witness that $>_{T_\psi}$ is ill founded is necessarily a proper subset of κ of type κ closed under ψ . Since this argument holds for an arbitrary $\psi \in W$, we can argue that W models that κ is Jónsson. This concludes the proof of both propositions. ■

The results that I presented suggest to investigate the following problem as the next step towards a solution of the above conjectures:

Problem 34 *Is it possible to prove the analogue of corollary 28 for sequences of size ω_1 ? I.e. is it possible to prove that if V models MM, W is an inner model and κ is a regular W -cardinal with $\kappa^+ = (\kappa^+)^W$, then $\text{cof}(\kappa) > \omega_1$ in V ?*

Assuming a positive answer to this problem, the next proposition shows that conjecture 1 cannot have a negative solution by means of set-forcing.

Proposition 35 *Assume $V \subseteq W$ are models of MM with the same ordinals of cofinality ω and ω_1 and that W is a set forcing extension of V . Then $[\text{Ord}]^{\omega_1} \subseteq V$.*

Proof: W is a set-generic extension of V by some P -generic filter. Thus P is a set and satisfies the $|P|^+$ -chain condition. Let $\kappa = |P|^+$. It is enough to show that $[\kappa]^{\omega_1} \subseteq V$. Now $S_\kappa^{\leq \omega_1} = (S_\kappa^{\leq \omega_1})^W$. By the κ -chain condition we get that every stationary subset of $S_\kappa^\omega \in V$ remains stationary in W . Now fix a partition $\{S_\alpha : \alpha < \kappa\}$ in V of S_κ^ω in κ -many disjoint stationary subset of κ . Then this is still a partition in stationary sets of W . Given any $g \in \kappa^{\leq \omega_1}$ find by MM in W an ordinal $\delta < \kappa$ of cofinality ω_1 such that $S_{g(\xi)}$ reflects on δ for all ξ in the domain of g . Let $C \in V$ be a club subset of δ of order type ω_1 such that $\alpha \in C$ iff there is ξ such that $\alpha \in S_{g(\xi)}$. Then $\text{Im}(g) \in V$ since:

$$\text{Im}(g) = \{\eta : \exists \alpha \in S_\eta \cap C\}.$$

We can conclude that also $g \in V$. ■

Thus also the following problem seems to be relevant to our discussion:

Problem 36 *Is it possible to produce two models $W \subseteq V$ of MM with the same cardinals, the same ordinals of cofinality ω and ω_1 and such that some stationary set of $(S_\kappa^\omega)^W$ is not anymore stationary in V , where κ is a regular cardinal of V ?*

References

- [1] U. Abraham and S. Todorćević. Partition Properties of ω_1 Compatible with CH. *Fundamenta Mathematicae*, 152:165–180, 1997.
- [2] B. Balcar, T. Jech, and T. Pazák. Complete ccc boolean algebras, the order sequential topology, and a problem of von neumann. *Bulletin of the London Mathematical Society*, 37:885–898, 2005.

- [3] R. E. Beaudoin. The proper forcing axiom and stationary set reflection. *Pacific Journal of Mathematics*, 149(1):13–24, 1991.
- [4] A. Caicedo and B. Veličković. The Mapping Reflection Principle. In preparation.
- [5] A. Caicedo and B. Veličković. Bounded proper forcing axiom and well orderings of the reals. *Mathematical Research Letters*, pages 393–408, 2006.
- [6] M. Gitik. The strength of the failure of SCH. *Annals of Pure and Applied Logics*, 51(3):215–240, 1991.
- [7] T. Jech. *Set theory, The Third Millennium Edition, Revised and Expanded*. Springer, 2002.
- [8] A. Kanamori and M. Magidor. The evolution of large cardinals axioms in set theory. In G. Müller and D. S. Scott, editors, *Higher Set theory, Proceedings Oberwolfach 1977*, pages 99–276. Springer-Verlag, 1978.
- [9] B. König. Forcing indestructibility of set-theoretic axioms. *Journal of Symbolic Logic*, 72:669–688, 2007.
- [10] B. König, P. Larson, J. T. Moore, and B. Veličković. Bounding the consistency strength of a five element linear basis. To appear in *Israel Journal of Mathematics*, 17 pages.
- [11] B. König and Y. Yoshinobu. Fragments of Martin’s Maximum in generic extensions. *Mathematical Logic Quarterly*, 50(3):297–302, 2004.
- [12] P. B. Larson. *The Stationary Tower: Notes on a Course by W. Hugh Woodin*. AMS, 2004.
- [13] T. Miyamoto. a memo. unpublished, 2 pages, 2006.
- [14] J. T. Moore. Set mapping reflection. *Journal of Mathematical Logic*, 5(1):87–97, 2005.
- [15] J. T. Moore. A five element basis for the uncountable linear orders. *Annals of Mathematics*, 163(2):669–688, 2006.
- [16] J. T. Moore. The Proper Forcing Axiom, Prikry forcing, and the Singular Cardinals Hypothesis. *Annals of Pure and Applied Logic*, 140(1-3):128–132, 2006.
- [17] E. Schimmerling. Combinatorial principles in the core model for one Woodin cardinal. *Annals of Pure and Applied Logic*, 74(2):153–201, 1995.
- [18] S. Shelah. *Cardinal Arithmetic*. Oxford University Press, 1994.
- [19] J. H. Silver. On the singular cardinal problem. *Proceedings of the International Congress of Mathematicians, Vancouver, B.C., 1974*, 1:265–268, 1975.

- [20] R. M. Solovay. Strongly compact cardinals and the GCH. *Proc. Sympos. Pure Math., Vol XXV, Univ. of California, Berkeley, Calif.*, pages 365–372, 1971.
- [21] S. Todorčević. A dichotomy for P -ideals of countable sets. *Fundamenta Mathematicae*, 166(3):251–267, 2000.
- [22] S. Todorčević. Coherent sequences. In M. Foreman, A. Kanamori, and M. Magidor, editors, *Handbook of Set Theory*. North Holland, to appear.
- [23] B. Veličković. Forcing axioms and stationary sets. *Advances in Mathematics*, 94(2):256–284, 1992.
- [24] B. Veličković. CCC Forcing and Splitting Reals. *Israel Journal of Mathematics*, 147:209–220, 2005.
- [25] M. Viale. The Proper Forcing Axiom and the Singular Cardinal Hypothesis. *Journal of Symbolic Logic*, 71(2):473–479, 2006.